IFIC/13-XXX

arXiv:1303.4685v1 [astro-ph.HE] 19 Mar 2013

Updated CMB, X- and gamma–ray constraints on majoron dark matter

Massimiliano Lattanzi,[1] Signe Riemer-Sørensen,[2] Mariam Tórtola,[3] and J. W. F. Valle[3, *]

[1] *Dipartimento di Fisica e Science della Terra, Università di Ferrara and INFN, sezione di Ferrara, Polo Scientifico e Tecnologico - Edificio C Via Saragat, 1, I-44122 Ferrara Italy*

[2] *School of Mathematics and Physics, University of Queensland, St Lucia, Brisbane 4072, Queensland, Australia*

[3] *AHEP Group, Instituto de Física Corpuscular – C.S.I.C./Universitat de València Campus de Paterna, Apt 22085, E–46071 València, Spain*

The majoron provides an attractive dark matter candidate, directly associated to the mechanism responsible for spontaneous neutrino mass generation within the standard model $\mathrm{SU(3)_c \otimes SU(2)_L \otimes U(1)_Y}$ framework. Here we update the cosmological and astrophysical constraints on majoron dark matter which come from the Cosmic Microwave Background (CMB) and a variety of X- and gamma–Ray observations.

PACS numbers: 98.80.Es, 98.80.Jk, 95.30.Sf

I. INTRODUCTION

It is by now established that only a small fraction (less than 20%) of the matter in the Universe is in the form of ordinary - i.e., baryonic - matter, while the rest is in the form of so-called "dark matter". The existence of dark matter is inferred by gravitational anomalies at very different scales, ranging from galactic scales to cluster scales, and all the way up to the cosmological scales. In particular, the recent 9-year data from the Wilkinson Microwave Anisotropy Probe (WMAP) [1, 2] have provided even stronger support to the six-parameter ΛCDM model, although small-scale experiments like ACT [3] and SPT [4] hint to interesting, albeit discordant, deviations from this simple picture, like e.g. the presence of other relativistic degrees of freedom, in addition to the standard model neutrinos, or deviations from ordinary gravity. The soon-to-be-released data from the Planck satellite will allow to measure the ΛCDM parameters with high precision, as well as to assess the significance of various extension to the simplest model.

If the ΛCDM model is certainly a phenomenological success, nevertheless it is puzzling from the theoretical point of view in many aspects. On one side, the nature of both dark matter and dark energy, that together make up for more than 95% of the total energy budget of the Universe, is still unknown. On the other side, the theory of inflation, that explains the formation of the primeval seeds for density fluctuations from which galaxies originate, is still waiting to be embedded in a more fundamental theory. All these puzzles hint, and possibly have their solution in, some physics beyond the standard model (SM) of particle physics, or maybe in some modification of general relativity.

Although its precise nature is still unknown, there is no shortage of candidates for the role of dark matter. One of the most widely studied candidates to date is the supersymmetric neutralino. Recent results from the Large Hadron Collider (LHC), however, have greatly reduced the available parameter space for supersymmetry, at least in its simplest minimal supergravity implementations [5], reducing, perhaps, the appeal of supersymmetric dark matter candidates. Other possibilities include the axion, sterile neutrinos, Kaluza-Klein dark matter, and many others.

The evidence for the existence of dark matter is very strong, but only limited to effects related to its gravitational interaction. The search for non-gravitational evidence of the dark matter continues, in the form of direct and indirect detection experiments, and by looking for dark matter production in accelerators like the LHC. A precise underpinning of the dark matter and determination of its properties can only come through a combination of these approaches.

If the dark matter has any connection to the world of SM particles, there will be astrophysical signals one can search for, in particular high energy photons from annihilating or decaying dark matter (see Ref. [6] for a recent review). The most studied scenarios are the broad spectrum annihilation signals from neutralinos, but the real smoking gun is line emission (either directly from the decay/annihilation [7, 8] or from internal bremsstrahlung [9, 10]), for which the spectral and spatial distribution is

---

[*] http://astroparticles.ific.uv.es/

not easily mimicked by astrophysical sources. The recent claim of line emission at $E_\gamma = 130\,\mathrm{GeV}$ in the Fermi data [11, 12] has spurred a renewed interest in emission line searches at high energy. However, the origin of the possible signal at $E_\gamma = 130\,\mathrm{GeV}$ is still unknown and requires further investigation.

It has been long suggested [13] that the origin of dark matter could be related to the origin of neutrino masses [14]. In fact, the smallness of neutrino masses, as compared to the other SM particles, is puzzling in itself. Most likely it is associated to the properties of the messenger states whose exchange is responsible for inducing them. This is the idea underlying the so-called seesaw mechanism [15–19], whose details remain fairly elusive. Especially appealing is the possibility that neutrino masses arise from the spontaneous violation of ungauged lepton number [20, 21]. The associated Nambu-Goldstone boson, the majoron, could acquire a mass from non-perturbative gravitational effects [22, 23], and play the role of the dark matter particle. In Ref. [24] the viability of the majoron as a dark matter particle was explored using the WMAP 3-year data and in Ref. [25] the possible X-ray signature associated to majoron decay was investigated. A specific theoretical model implementing the seesaw mechanism and an A_4 flavour symmetry was described in [26].

In this paper, we update our previous constraints in the light of the more recent cosmological and astrophysical data. Regarding cosmology we take into account the most recent WMAP 9-year data [1, 2]. On the astrophysical front we include emission line searches on the entire range of photon energies between $0.07\,\mathrm{keV}$ and $200\,\mathrm{GeV}$ from *Chandra* X-ray Observatory, X-ray Multi-Mirror Mission - Newton (*XMM*), High Energy Astronomy Observatory Program (*HEAO*), INTErnational Gamma-Ray Astrophysics Laboratory (*INTEGRAL*), Compton Gamma Ray Observatory (*CGRO*), and the Fermi Gamma-ray Space Telescope.

The paper is organized as follows. In Sec II, we briefly recall the basics of majoron physics.. In Secs. III and IV, we derive observational constraints on the majoron decay to neutrinos and photons, respectively, and we compare them to the predictions of a general seesaw model. Finally, in Sec. V we draw our conclusions.

II. SEESAW MAJORON PHYSICS

The basic idea is that the lepton number symmetry of the standard $\mathrm{SU(3)_c \otimes SU(2)_L \otimes U(1)_Y}$ model is promoted to a spontaneously broken symmetry [20, 21]. This requires the presence of a lepton-number-carrying complex scalar singlet coupling to the singlet neutrinos ν^c_L, as follows,

$$\lambda \sigma {\nu^c_L}^T \sigma_2 \nu^c_L + H.c. \tag{1}$$

This term provides the large mass term in the seesaw mass matrix

$$\mathcal{M}_\nu = \begin{bmatrix} Y_3 v_3 & Y_\nu v_2 \\ {Y_\nu}^T v_2 & Y_1 v_1 \end{bmatrix} \tag{2}$$

in the basis ν_L, ν^c_L of "left" and "right"-handed neutrinos. The model is characterized by singlet, doublet and triplet Higgs scalars whose vacuum expectation values (vevs) are arranged to satisfy $v_1 \gg v_2 \gg v_3$ obeying a simple vev seesaw relation of the type

$$v_3 v_1 \sim {v_2}^2. \tag{3}$$

The vev v_1 drives lepton number violation and induces also a small but nonzero v_3 while ${v_2}^2$ is fixed by the masses of the weak gauge bosons, the W and the Z. Note that the vev seesaw condition implies that the triplet vev $v_3 \to 0$ as the singlet vev $v_1 \to \infty$. The three vevs determine all entries in the seesaw neutrino mass matrix. Regarding the Yukawa couplings, Y_ν is an arbitrary flavour matrix, while Y_3 and Y_1 are symmetric. The effective light neutrino mass obtained by perturbative diagonalization of Eq. (2) is of the form

$$m_\nu \simeq Y_3 v_3 - Y_\nu {Y_1}^{-1} {Y_\nu}^T \frac{{v_2}^2}{v_1} \tag{4}$$

Together with Eq. (3) this summarizes the essence of the seesaw mechanism.

In order to identify which combination of Higgs fields gives the majoron, one may write the scalar potential explicitly, minimize it, and determine the resulting scalar mass matrices. However, one can do this simply by exploiting the invariance properties of the Higgs potential V [21]. The result is proportional to the combination

$$J \propto v_3 {v_2}^2 \Im(\Delta^0) - 2 v_2 {v_3}^2 \Im(\Phi^0) + v_1({v_2}^2 + 4{v_3}^2)\Im(\sigma) \tag{5}$$

up to normalization (note $\Im$ denotes imaginary parts). We remark the presence of the quartic lepton-number-conserving term

$$\Phi^\dagger \Delta \tau_2 \Phi^* \sigma^* + H.c. \tag{6}$$

in the scalar potential. Here τ_2 is the weak isospin Pauli matrix, and $v_2 \equiv \langle\Phi\rangle$, $v_1 \equiv \langle\sigma\rangle$, $v_3 \equiv \langle\Delta\rangle$. This term illustrates the need for mixing among neutral fields belonging to all three Higgs multiplets in the expression for the majoron, Eq. (5). This implies that the majoron has an explicit coupling to two photons that leads to the possibility of indirect detection of majoron dark matter by searching for the corresponding high energy photons [25], which we treat in Sec. IV.

We now turn to the form of the couplings of the majoron within the above seesaw scheme, characterized by spontaneous lepton number violation in the presence of singlet, doublet and triplet Higgs scalars. Again one can derive the form of the couplings of the majoron using only the symmetry properties, as described in Ref. [21],

$$\mathcal{L}_{\mathrm{Yuk}} = \frac{iJ}{2}\sum_{ij}\nu_i^T g_{ij}\sigma_2\nu_j + H.c. \tag{7}$$

The result is a perturbative expansion for the majoron couplings

$$g_{ij} = -\frac{m_i}{v_1}\delta_{ij} + \dots \tag{8}$$

where the dots ... denote higher order terms. One sees that, to first approximation, the majoron couples to the light mass-eigenstate neutrinos inversely proportional to the lepton number violation scale $v_1 \equiv \langle\sigma\rangle$ and proportionally to their mass. With this we can compute the dark matter majoron decay rate to neutrinos as

$$\Gamma_{J\nu\nu} = \frac{m_J}{32\pi}\frac{\sum_i (m_i^\nu)^2}{2v_1^2}\ . \tag{9}$$

Moreover, there is a sub-leading majoron decay mode to gammas. Within the general seesaw model this decay is induced at the loop level, resulting in [25]

$$\Gamma_{J\gamma\gamma} = \frac{\alpha^2 m_J^3}{64\pi^3}\left|\sum_f N_f Q_f^2 \frac{2v_3^2}{v_2^2 v_1}(-2T_3^f)\frac{m_J^2}{12m_f^2}\right|^2 , \tag{10}$$

where N_f, Q_f, T_3^f and m_f denote respectively the color factor, electric charge, weak isospin and mass of the SM electrically charged fermions f. We note that this formula is an approximation valid for $m_J \ll m_f$; however we will always use the exact formula in the actual calculations.

The decay of the majoron dark matter to neutrinos provides the most essential and model–independent feature of the majoron dark matter scenario, namely, it is a decaying dark matter model where the majoron decays mainly to neutrinos, a mode that is constrained from the CMB observations, as we discuss in Sec. III.

III. CMB CONSTRAINTS ON THE INVISIBLE DECAY $J \to \nu\nu$

We start by deriving constraints on the majoron properties from CMB anisotropy data. The majoron differs from most dark matter candidates in that it is unstable, since it must decay to neutrinos, as seen in Eq. (9), although it obviously has to be long-lived enough to be the dark matter.

In order to investigate the observable effects of majoron decay on the CMB, the Boltzmann equation describing the phase-space evolution of DM particles must be modified accordingly, as shown e.g. in Ref. [27], both at the background and at the perturbation level. The main effect of the late DM decay to invisible relativistic particles is the increase of the late integrated Sachs-Wolfe (LISW), caused by the presence of an extra radiation component at small redshifts. This is reflected in an increased amount of power at the largest angular scales (i.e., small multipoles). Too large a decay rate would produce too much radiation and too much large-scale power, and thus be at variance with observations. In Ref. [24] two of us have used this effect to constrain the majoron lifetime. However, we did not take properly into account the effect of majoron decay on the age of the Universe; this lead to an underestimate of the upper limit of the majoron decay rate. We have now corrected this.

We use a modified version of CAMB [28], taking into account the finite lifetime of the majoron, to evolve the cosmological perturbations and compute the anisotropy spectrum of the CMB for given values of the cosmological parameters. In order to compute credible intervals and sample the posterior distributions for the parameters of the model, given some data, we use the Metropolis-Hastings algorithm as implemented in CosmoMC [29] (interfaced with our modified version of CAMB). We take uniform priors on all the parameters of the model, namely the present density parameters $\Omega_{\mathrm{b}}h^2$ and $\Omega_{\mathrm{dm}}h^2$ of baryons and dark matter respectively, the Hubble constant H_0, the optical depth to recombination τ_{rec}, the spectral index n_s and amplitude A_s (evaluated at the pivot scale $k_0 = 0.002\ \mathrm{Mpc}^{-1}$) of the spectrum of primordial scalar fluctuations, and the decay rate $\Gamma_{J\to\nu\bar{\nu}}$ of majoron to neutrinos. We marginalize over the amplitude of the contamination from the Sunyaev-Zel'dovich signal.

We perform our analysis using the most recent WMAP 9-year temperature and polarization data [1, 2]. In particular, in temperature we include data up to $\ell_{\max} =$



FIG. 1. Two-dimensional WMAP-9 constraints on the majoron dark matter present density and decay rate to neutrinos. The light (dark) shaded regions correspond to 68% (95%) credible intervals.

1200. We use the latest (V5) version of their likelihood code, publicly avaiilable at the lambda website[1].

For the purpose of the present analysis, we are mainly interested in the limits on the decay width of majoron to neutrinos, $\Gamma_{J\to\nu\nu}$. We get the following upper limit at 95% C.L.

$$\Gamma_{J\to\nu\nu} \leq 6.4\times 10^{-19}\,{\rm s}^{-1}\,. \tag{11}$$

resulting in a lower limit to the majoron lifetime $\tau_J \geq$ 50 Gyr, roughly four times the age of the Universe. We have also checked that this limit does not substantially change when we allow for the possibility of extra degrees of freedom at the time of recombination.

In Fig. 1 we show 68% and 95% credible intervals in the ($\Gamma_{J\to\nu\nu}$, $\Omega_{\rm dm}h^2$) plane. There is an evident anticorrelation that is explained by the fact, already discussed in Ref. [24], that the CMB is mainly sensitive to the amount of dark matter at early times (through the height of the first peak in the anisotropy spectrum). For a given amount of dark matter in the early Universe, a larger decay rate would result in a smaller present amount of dark matter.



FIG. 2. 3σ line emission constraints on the decay rate into two mono-energetic photons. Applies to all dark matter candidates with this signature. The constraints are taken from yellow [30], orange [31] (conservatively rescaled by a factor of two due to mass estimate uncertainties as recommended in [32]), red [33], grey [34], purple [35], blue [36], cyan [37], green [38].

IV. X- AND γ-RAY CONSTRAINTS ON THE PHOTON DECAY $J\to\gamma\gamma$

One of the most interesting features of spontaneous lepton number violation within the general $\mathrm{SU(3)_c\otimes SU(2)_L\otimes U(1)_Y}$ seesaw model is that the neutrino decay mode in Eq. (9) is accompanied by a two-photon mode, Eq. (10), as a result of the Eq. (5). This is constrained by a number of astrophysical observations.

A. Existing constraints

In Fig. 2 we plot the emission line constraints over the wide range of photon energies of 0.07 keV to 200 GeV.

The very soft X-ray emission is covered by *Chandra* Low Energy Transmission Grating (LETG) observations of NGC3227 (0.07 − 4.1 keV) [30] and a rocket borne light cryogenic spectrometer (0.25 − 1.1 keV) [31, 39].

The 0.3 − 12 keV range is well covered with constraints from various objects observed with the *Chandra* and *XMM* X-ray telescopes [33–36, 40–48] In Fig. 2 we have chosen the strongest robust constraints from *XMM* observations of the Milky Way and M31 [33] and from *Chandra* observations of the Draco dwarf galaxy [34][2].

---

[1] http://lambda.gsfc.nasa.gov/

The diffuse X-ray background observed with HEAO was searched for line emission by Ref. [35] over the range $3 - 48\,\text{keV}$. Line emission constraints have been derived from INTEGRAL SPI observations of the soft γ-ray background ($20\,\text{keV} - 7\,\text{MeV}$) [36]. For energies above those covered by INTEGRAL the constraints are two orders of magnitude worse as this range is only covered by a combination of the rather old COMPTEL and EGRET instruments onboard the CGRO. However, line emission constraints have been derived up to $100\,\text{GeV}$ [37]. The most recent flagship for γ-ray searches is the Fermi γ-ray Space Telescope, for which line emission searches have been performed for the range of $7 - 200\,\text{GeV}$ [38].

B. Future sensitivities

The constraints in Fig. 2 can be improved by increasing the statistics or the spectral resolution. Increasing statistics (either by exposure time or by sensitivity) improves the constraints as $\Gamma_\gamma^{\text{new}} = \sqrt{N_\gamma^{\text{ex}}/N_\gamma^{\text{new}}}\Gamma_\gamma^{\text{ex}}$, where $N_\gamma^{\text{ex, new}}$ are the existing and new total of photons per bin (assuming both source and background counts increase by the same amount). Increasing the spectral resolution improves the constraints directly as $\Gamma_\gamma^{\text{new}} = E_{\text{FWHM}}^{\text{new}}/E_{\text{FWHM}}^{\text{ex}}\Gamma_\gamma^{\text{ex}}$ and is consequently preferable but also technically more challenging.

C. Model comparison

We now compare the observational constraints obtained in the previous section to the predictions of different realizations of a general majoron seesaw model. In particular, we perform a random scan over the Yukawa matrices and vevs that characterize the seesaw mass matrix $\mathcal{M}_\nu$ in Eq. (2). For each point in the parameter space we evaluate the effective light neutrino mass matrix and the Majoron decay rate to neutrinos following Eqs. (4) and (9), respectively. We then choose, among all possible realizations, those that are in agreement with current neutrino oscillation data [53] as well as with the neutrino bound in Eq. (11). Finally we compute the corresponding decay rate to photons, as described in Sec. II.



FIG. 3. The line emission constraints from Figure 2 (grey) compared to model predictions (colored dots), for different values of the triplet vev v_3.

We show the results of our scan in parameter space in Fig. 3, together with the constraints already shown in Fig. 2. It is clearly visible that the $J \to \gamma\gamma$ constraints from line emission searches already begin to cut the parameter space for realistic models. This happens in particular for models with v_3 larger than a few MeVs. However, models with lower values of the triplet vev predict a photon flux that below the observational limits by at least 6 orders of magnitude.

V. CONCLUSIONS

We have updated previous constraints on the parameters of the majoron dark matter model using the most recent CMB, X- and γ–ray observations. From the CMB, we have derived an upper limit on the rate of the invisible decay of the dark matter particle, namely, in the framework of the model under consideration, on the majoron decay to neutrinos. Translated in terms of the particle lifetime, this constraints the majoron lifetime to be larger than 50 Gyrs.

---

[2] Some analyses have claimed stronger constraints, but later found to be too optimistic. Ref. [49] underestimated the flux by two orders of magnitude [40, 42]. According to Ref. [33] the mass was overestimated in Ref. [50] leading to too restrictive constraints. The constraints in Ref. [51] might be too restrictive due to the choice of source profile [36], and the spectral resolution appears overestimated in Ref. [52]

The majoron also possesses a subleading decay mode to two photons, that can be constrained by astrophysical observations in the X and γ regions. We have compared these limits to the theoretical predictions corresponding to different values for the parameters of the underlying particle physics model. We have found that the observational constraints already exclude part of the parameter space for models in which the vev of the triplet v_3 is larger than a few MeVs. On the other hand, for smaller values of v_3, the current limits need to be improved by at least 6 orders of magnitude before the allowed region in parameter space can be reduced.

VI. ACKNOWLEDGMENTS

Work supported by MINECO grants FPA2011-22975 and MULTIDARK Consolider CSD2009-00064, by Prometeo/2009/091 (Gen. Valenciana), by EU ITN UNILHC PITN-GA-2009-237920. The work of M.L. has been supported by Ministero dell'Istruzione, dell'Università e della Ricerca (MIUR) through the PRIN grant "Galactic and extragalactic polarized microwave emission" (contract number PRIN 2009XZ54H2-002). The work of M.T. is supported by CSIC under the JAE-Doc programme, co-funded by the European Social Fund.

[1] G. Hinshaw, D. Larson, E. Komatsu, D. Spergel, C. Bennett, *et al.*, (2012), arXiv:1212.5226 [astro-ph.CO].

[2] C. Bennett, D. Larson, J. Weiland, N. Jarosik, G. Hinshaw, *et al.*, (2012), arXiv:1212.5225 [astro-ph.CO].

[3] J. L. Sievers, R. A. Hlozek, M. R. Nolta, V. Acquaviva, G. E. Addison, *et al.*, (2013), arXiv:1301.0824 [astro-ph.CO].

[4] Z. Hou, C. Reichardt, K. Story, B. Follin, R. Keisler, *et al.*, (2012), arXiv:1212.6267 [astro-ph.CO].

[5] S. Caron (ATLAS collaboration), (2011), arXiv:1106.1009 [hep-ex].

[6] T. Bringmann and C. Weniger, Physics of the Dark Universe **1**, 194 (2012), arXiv:1208.5481 [hep-ph].

[7] L. Bergström and P. Ullio, Nuclear Physics B **504**, 27 (1997), arXiv:hep-ph/9706232.

[8] Z. Bern, P. Gondolo, and M. Perelstein, Physics Letters B **411**, 86 (1997), arXiv:hep-ph/9706538.

[9] J. F. Beacom, N. F. Bell, and G. Bertone, Physical Review Letters **94**, 171301 (2005), arXiv:astro-ph/0409403.

[10] T. Bringmann, L. Bergström, and J. Edsjö, Journal of High Energy Physics **1**, 049 (2008), arXiv:0710.3169 [hep-ph].

[11] T. Bringmann, X. Huang, A. Ibarra, S. Vogl, and C. Weniger, JCAP **7**, 054 (2012), arXiv:1203.1312 [hep-ph].

[12] C. Weniger, JCAP **8**, 007 (2012), arXiv:1204.2797 [hep-ph].

[13] G. Gelmini, D. N. Schramm, and J. W. F. Valle, Phys. Lett. **B146**, 311 (1984).

[14] V. Berezinsky and J. W. F. Valle, Phys. Lett. **B318**, 360 (1993), hep-ph/9309214.

[15] M. Gell-Mann, P. Ramond, and R. Slansky, (1979), print-80-0576 (CERN).

[16] T. Yanagida, (KEK lectures, 1979), ed. O. Sawada and A. Sugamoto (KEK, 1979).

[17] R. N. Mohapatra and G. Senjanovic, Phys. Rev. Lett. **44**, 91 (1980).

[18] J. Schechter and J. W. F. Valle, Phys. Rev. **D22**, 2227 (1980).

[19] G. Lazarides, Q. Shafi, and C. Wetterich, Nucl. Phys. **B181**, 287 (1981).

[20] Y. Chikashige, R. N. Mohapatra, and R. D. Peccei, Phys. Lett. **B98**, 265 (1981).

[21] J. Schechter and J. W. F. Valle, Phys. Rev. **D25**, 774 (1982).

[22] S. R. Coleman, Nucl. Phys. **B310**, 643 (1988).

[23] R. Kallosh, A. D. Linde, D. A. Linde, and L. Susskind, Phys. Rev. **D52**, 912 (1995), arXiv:hep-th/9502069.

[24] M. Lattanzi and J. W. F. Valle, Phys. Rev. Lett. **99**, 121301 (2007), arXiv:0705.2406 [astro-ph].

[25] F. Bazzocchi *et al.*, JCAP **0808**, 013 (2008), arXiv:0805.2372 [astro-ph].

[26] J. Esteves, F. Joaquim, A. Joshipura, J. Romao, M. Tortola, *et al.*, Phys.Rev. **D82**, 073008 (2010), arXiv:1007.0898 [hep-ph].

[27] M. Kaplinghat, R. E. Lopez, S. Dodelson, and R. J. Scherrer, Phys. Rev. **D60**, 123508 (1999), astro-ph/9907388.

[28] A. Lewis, A. Challinor, and A. Lasenby, Astrophys. J. **538**, 473 (2000), astro-ph/9911177.

[29] A. Lewis and S. Bridle, Phys. Rev. **D66**, 103511 (2002), astro-ph/0205436.

[30] F. Bazzocchi, M. Lattanzi, S. Riemer-Sorensen, and J. W. F. Valle, JCAP **8**, 013 (2008), arXiv:0805.2372.

[31] A. Boyarsky, J.-W. den Herder, A. Neronov, and O. Ruchayskiy, Astroparticle Physics **28**, 303 (2007), arXiv:astro-ph/0612219.

[32] A. Boyarsky, O. Ruchayskiy, and M. Shaposhnikov, Annual Review of Nuclear and Particle Science **59**, 191 (2009), arXiv:0901.0011 [hep-ph].

[33] A. Boyarsky, D. Iakubovskyi, O. Ruchayskiy, and V. Savchenko, MNRAS **387**, 1361 (2008), arXiv:0709.2301.

[34] S. Riemer-Sorensen and S. H. Hansen, ArXiv e-prints (2009), arXiv:0901.2569 [astro-ph.CO].

[35] A. Boyarsky, A. Neronov, O. Ruchayskiy, and M. Shaposhnikov, MNRAS **370**, 213 (2006), arXiv:astro-ph/0512509.

[36] A. Boyarsky, D. Malyshev, A. Neronov, and O. Ruchayskiy, MNRAS **387**, 1345 (2008), arXiv:0710.4922.

[37] H. Yüksel and M. D. Kistler, Phys. Rev. D **78**, 023502 (2008), arXiv:0711.2906.

[38] M. Ackermann, M. Ajello, A. Albert, L. Baldini, G. Barbiellini, K. Bechtol, R. Bellazzini, B. Berenji, R. D. Blandford, E. D. Bloom, E. Bonamente, A. W. Borgland, M. Brigida, R. Buehler, S. Buson, G. A. Caliandro, R. A. Cameron, P. A. Caraveo, J. M. Casandjian, C. Cecchi, E. Charles, A. Chekhtman, J. Chiang, S. Ciprini, R. Claus, J. Cohen-Tanugi, J. Conrad, F. D'Ammando, F. de Palma, C. D. Dermer, E. do Couto e Silva, P. S. Drell, A. Drlica-Wagner, Y. Edmonds, R. Essig, C. Favuzzi, S. J. Fegan, W. B. Focke, Y. Fukazawa, S. Funk, P. Fusco, F. Gargano, D. Gasparrini, S. Germani, N. Giglietto, F. Giordano, M. Giroletti, T. Glanzman, G. Godfrey, I. A. Grenier, S. Guiriec, M. Gustafsson, D. Hadasch, M. Hayashida, D. Horan, R. E. Hughes, T. Kamae, J. Knödlseder, M. Kuss, J. Lande, A. M. Lionetto, M. Llena Garde, F. Longo, F. Loparco, M. N. Lovellette, P. Lubrano, M. N. Mazziotta, P. F. Michelson, W. Mitthumsiri, T. Mizuno, A. A. Moiseev, C. Monte, M. E. Monzani, A. Morselli, I. V. Moskalenko, S. Murgia, M. Naumann-Godo, J. P. Norris, E. Nuss, T. Ohsugi, A. Okumura, E. Orlando, J. F. Ormes, D. Paneque, J. H. Panetta, M. Pesce-Rollins, F. Piron, G. Pivato, T. A. Porter, D. Prokhorov, S. Rainò, R. Rando, M. Razzano, O. Reimer, M. Roth, C. Sbarra, J. D. Scargle, C. Sgrò, E. J. Siskind, A. Snyder, P. Spinelli, D. J. Suson, H. Takahashi, T. Tanaka, J. G. Thayer, J. B. Thayer, L. Tibaldo, M. Tinivella, D. F. Torres, G. Tosti, E. Troja, J. Vandenbroucke, V. Vasileiou, G. Vianello, V. Vitale, A. P. Waite, B. L. Winer, K. S. Wood, Z. Yang, and S. Zimmer, Phys. Rev. D **86**, 022002 (2012), arXiv:1205.2739 [astro-ph.HE].

[39] D. McCammon, R. Almy, E. Apodaca, W. Bergmann Tiest, W. Cui, S. Deiker, M. Galeazzi, M. Juda, A. Lesser, T. Mihara, J. P. Morgenthaler, W. T. Sanders, J. Zhang, E. Figueroa-Feliciano, R. L. Kelley, S. H. Moseley, R. F. Mushotzky, F. S. Porter, C. K. Stahle, and A. E. Szymkowiak, Astrophys. J. **576**, 188 (2002), arXiv:astro-ph/0205012.

[40] A. Boyarsky, A. Neronov, O. Ruchayskiy, M. Shaposhnikov, and I. Tkachev, Physical Review Letters **97**, 261302 (2006), arXiv:astro-ph/0603660.

[41] S. Riemer-Sørensen, S. H. Hansen, and K. Pedersen, ApJL **644**, L33 (2006), arXiv:astro-ph/0603661.

[42] K. Abazajian and S. M. Koushiappas, Phys. Rev. D **74**, 023527 (2006), arXiv:astro-ph/0605271.

[43] A. Boyarsky, J. Nevalainen, and O. Ruchayskiy, A&A **471**, 51 (2007), arXiv:astro-ph/0610961.

[44] S. Riemer-Sorensen, K. Pedersen, S. H. Hansen, and H. Dahle, Phys. Rev. D **76**, 043524 (2007), arXiv:astro-ph/0610034.

[45] M. Loewenstein, A. Kusenko, and P. L. Biermann, Astrophys. J. **700**, 426 (2009), arXiv:0812.2710.

[46] M. Loewenstein and A. Kusenko, Astrophys. J. **714**, 652 (2010), arXiv:0912.0552 [astro-ph.HE].

[47] M. Loewenstein and A. Kusenko, Astrophys. J. **751**, 82 (2012), arXiv:1203.5229 [astro-ph.CO].

[48] E. Borriello, M. Paolillo, G. Miele, G. Longo, and R. Owen, MNRAS **425**, 1628 (2012), arXiv:1109.5943 [astro-ph.GA].

[49] K. Abazajian, G. M. Fuller, and W. H. Tucker, Astrophys. J. **562**, 593 (2001), arXiv:astro-ph/0106002.

[50] C. R. Watson, J. F. Beacom, H. Yüksel, and T. P. Walker, Phys. Rev. D **74**, 033009 (2006), arXiv:astro-ph/0605424.

[51] H. Yüksel, J. F. Beacom, and C. R. Watson, Physical Review Letters **101**, 121301 (2008), arXiv:0706.4084.

[52] C. R. Watson, Z. Li, and N. K. Polley, JCAP **3**, 018 (2012), arXiv:1111.4217 [astro-ph.CO].

[53] D. Forero, M. Tortola, and J. W. F. Valle, Phys.Rev. **D86**, 073012 (2012), this updates New J.Phys. 13 (2011) 109401 and New J.Phys. 13 (2011) 063004 by including the data presented at Neutrino 2012., arXiv:arXiv:1205.4018 [hep-ph].